arXiv:1204.0412v3 [hep-ph] 3 Oct 2012

Spectrum of the excited N^* and Δ^* baryons in a relativistic chiral quark model

E.M. Tursunov
Institute of Nuclear Physics,
Uzbekistan Academy of Sciences,
100214, Ulugbek, Tashkent, Uzbekistan

(Dated: March 2, 2013)

The spectrum of the SU(2) flavor baryons is studied in the frame of a relativistic chiral quark potential model based on the one-pion and one-gluon exchange mechanisms. It is argued that the N^* and Δ^* resonances appearing in the πN scattering data and strongly coupled to the πN channel are identified with the orbital configurations $(1S_{1/2})^2(nlj)$ with a single valence quark in the excited state (nlj). Based on obtained selection rules we show that it is possible to construct a schematic periodic table of baryon resonances, consistent with the experimental data.

The numerical estimations for the energy positions of the Nucleon and Delta baryons (up to and including F-wave N^* and Δ^* resonances), obtained within the field-theoretical framework by using time ordered perturbation theory, yield an overall good description of the experimental data at a level of the relativized Constituent quark model of S. Capstick and W. Roberts without any fitting parameters. The only free parameter of the linear confinement potential was fitted previously by Th. Gutsche to reproduce the axial charge of the Nucleon. However, Nucleon ground state and some of the radially excited baryon resonances (including Roper) are overestimated. Contrary, all the orbitally excited N^* and some of the Δ^* resonances are underestimated. At the same time, the remaining four Nucleon and five Δ states including $\Delta(1232)$ are well reproduced.

At higher energies, where the experimental data are poor, we can extend our model schematically and predict an existence of seven N^* and four Δ^* new states with larger spin values.

PACS numbers: 11.10.Ef,12.39.Fe,12.39.Ki

I. INTRODUCTION

It was believed during the long period that the baryon spectrum can be described with a good accuracy in the Constituent Quark Models (CQM) based either on the Goldstone-boson exchange (GBE) [1], or one-gluon exchange (OGE) [2, 3] or (and) instanton induced exchange (IIE) [4] mechanisms between non-relativistic constituent quarks. However, there are still some serious problems within this framework which cannot be avoided until now. The most important issue is the problem of "missing resonances": the CQMs predicted too many states even at low energies which are not observed at the experimental facilities [5–7]. The situation is so serious, that "a modern view questions the usefulness of quarks to describe the nucleon excitation spectrum" [7].

Fortunately, there are also strong optimistic views on the problem. The recent review [6] shows all the difficulties in baryon spectroscopy and concludes that the all photo-, pion- and hadron-induced reactions will be important to understand the excited baryon spectroscopy. From the theoretical side, essential developments are being done in the Lattice QCD [8, 9], Dyson-Schwinger equations [10], effective field-theoretical methods [11–13] and in the theoretical coupled channel approaches to meson-baryon scattering within the Juelich [14] and Sato-Lee [15] models.

Although the relativistic chiral quark models [16–20] have been developed during the long period, there are no study of the excited baryon spectrum within these approaches in the literature. Since the chiral symmetry plays a special role for the hadron structure, it is important to probe these models in the excited baryon spectroscopy.

In [21–23] we have developed a relativistic chiral quark model for the lower excitation spectrum of the nucleon and delta. The splitting of the Roper resonance from the N(939) was reproduced with a reasonable accuracy. The model was tested firstly in Ref. [24] for the study of the Nucleon charge form factors, then in Ref. [25, 26] for the study of the nucleon properties such as mass, charge radius, magnetic moment, axial charge and reasonable agreement with the experimental values was obtained.

The model is based on an effective chiral Lagrangian. Quark wave function is obtained from the solution of the Dirac equation with a Cornell type potential containing a linear confining term and a Coulomb part due-to short range gluon field correlations. All the model parameters of the model, except one are fixed from the Lattice study of previous authors [27, 28]. The only free parameter of the model is the so-called "mass term" in the confinement potential, which was fitted in Ref.[26] to reproduce the axial charge of the Nucleon. The calculations are done at one loop or at order of accuracy $o(1/f_\pi^2, \alpha_s)$.

The aim of present paper is to extend the relativistic chiral quark model to the higher excitation spectrum of SU(2) flavor baryons. Firstly we want to check, whether the relativistic chiral quark model can help to understand the systematics of the excited Nucleon and Delta states and an orbital structure of each baryon state. Based on selection rules obtained from one-pion and one gluon exchange mechanisms between valence quarks, below we will show that it is possible to construct a periodic table, where each excited Nucleon or Delta state can be identified with an orbital configuration $(1S)^2(nlj)$ with a single radially or/and orbitally excited valence quark.

Secondly we will estimate the excited Nucleon and Delta spectrum in the present model with taking into account second-order perturbative corrections due-to the pion and color-magnetic gluon fields and compare with the experimental data.

The relevant suggestion is that the results of our study can be reproduced in any chiral quark model describing the baryons as bound states of three valence quarks with a Dirac two-component structure and surrounded by the cloud of π-mesons, as required by the chiral symmetry [29].

In Section 2 we give the main formalism of the model. The numerical results are presented in Section 3, and final conclusions are given in Section 4.

II. MODEL

A. Basis formalism

The effective Lagrangian of the model $\mathcal{L}(x)$ (see [25, 30]) contains the quark core part $\mathcal{L}_Q(x)$, the quark-pion $\mathcal{L}_I^{(q\pi)}(x)$ and the quark-gluon $\mathcal{L}_I^{(qg)}(x)$ interaction terms, and the kinetic parts for the pion $\mathcal{L}_\pi(x)$ and gluon $\mathcal{L}_g(x)$ fields:

$$\begin{aligned}\mathcal{L}(x) &= \mathcal{L}_Q(x) + \mathcal{L}_I^{(q\pi)}(x) + \mathcal{L}_I^{(qg)}(x) + \mathcal{L}_\pi(x) + \mathcal{L}_g(x)\\ &= \bar{\psi}(x)[i\,\partial\!\!\!/ - S(r) - \gamma^0 V(r)]\psi(x) - 1/f_\pi \bar{\psi}[S(r) i\gamma^5 \tau^i \phi_i]\psi - \\ &- g_s \bar{\psi} A^a_\mu \gamma^\mu \frac{\lambda^a}{2}\psi + \frac{1}{2}(\partial_\mu \phi_i)^2 - \frac{1}{2} m_\pi^2 \phi_i^2 - \frac{1}{4} G^a_{\mu\nu} G^{\mu\nu}_a . \end{aligned} \quad (1)$$

Here, $\psi(x)$, $\phi_i, i = 1,2,3$ and A^a_μ are the quark, pion and gluon fields, respectively. The matrices $\tau^i (i = 1,2,3)$ and $\lambda^a (a = 1, ..., 8)$ are the isospin and color matrices, correspondingly. The pion decay constant f_π =93 MeV. In the model, the chiral symmetry violated through the quark confinement mechanism is restored with the help of the linearized σ-model. The mass term for the pion field is introduced in order to satisfy the PCAC theorem [31], which is consistent with the Goldberger-Treiman relation.

We use the Cornell type potential in the Dirac equation for the single quark states in accordance with the Lattice QCD theory. The scalar part of the static confinement potential is given by

$$S(r) = cr + m \quad (2)$$

where c and m are constants. The strength parameter c of the confinement potential is defined from the Lattice study [27], while m is the only free parameter of the model which can be fitted to reproduce the axial charge g_A of the proton (and the πNN coupling constant via the Goldberger-Treiman relation).

At short distances, transverse fluctuations of the string are dominating [32], with an indication that they transform like the time component of the Lorentz vector. They are given by a Coulomb type vector potential (the so called Luscher term) as

$$V(r) = -\alpha/r \quad (3)$$

where $\alpha = \pi/12$ is defined from the QCD Lattice study [28].

The quark fields are obtained from solving the Dirac equation with the corresponding scalar plus vector potentials

$$[i\gamma^\mu \partial_\mu - S(r) - \gamma^0 V(r)]\psi(x) = 0 \quad (4)$$

The respective positive and negative energy eigenstates as solutions to the Dirac equation with a spherically symmetric mean field, are given in a general form as

$$u_\alpha(x) = \begin{pmatrix} g^+_{N\kappa}(r) \\ -i f^+_{N\kappa}(r)\, \vec{\sigma}\hat{\vec{x}} \end{pmatrix} \mathcal{Y}_\kappa^{m_j}(\hat{\vec{x}})\, \chi_{m_t}\, \chi_{m_c}\, exp(-iE_\alpha t) \quad (5)$$

$$v_\beta(x) = \begin{pmatrix} g^-_{N\kappa}(r) \\ -i f^-_{N\kappa}(r)\, \vec{\sigma}\hat{\vec{x}} \end{pmatrix} \mathcal{Y}_\kappa^{m_j}(\hat{\vec{x}})\, \chi_{m_t}\, \chi_{m_c}\, exp(+iE_\beta t) \quad (6)$$

The quark and anti-quark eigenstates u and v are labeled by the radial, angular, azimuthal, isospin and color quantum numbers N, κ, m_j, m_t and m_c, which are collectively denoted by α and β, respectively. The spin-angular part of the quark field operators

$$\mathcal{Y}_{\kappa}^{m_j}(\hat{\vec{x}}) \;=\; [Y_l(\hat{\vec{x}}) \otimes \chi_{1/2}]_{jm_j} \; j = |\kappa| - 1/2. \tag{7}$$

The quark fields ψ are expanded over the basis of positive and negative energy eigenstates as

$$\psi(x) = \sum_{\alpha} u_{\alpha}(x) b_{\alpha} + \sum_{\beta} v_{\beta}(x) d_{\beta}^{\dagger}. \tag{8}$$

The expansion coefficients b_α and $d_\beta^\dagger$ are operators, which annihilate a quark and create an anti-quark in the orbits α and β, respectively.

The free pion field operator is expanded over plane wave solutions as

$$\phi_j(x) = (2\pi)^{-3/2} \int \frac{d^3k}{(2\omega_k)^{1/2}} [a_{j\mathbf{k}} exp(-ikx) + a_{j\mathbf{k}}^{\dagger} exp(ikx)] \tag{9}$$

with the usual destruction and creation operators $a_{j\mathbf{k}}$ and $a_{j\mathbf{k}}^\dagger$ respectively. The pion energy is defined as $\omega_k \;=\; \sqrt{k^2 + m_\pi^2}$. The free gluon field operators is expanded in the same way.

In denoting the three-quark vacuum state by $|0>$, the corresponding noninteracting many-body quark Green's function (propagator) of the quark field is given by the customary vacuum Feynman propagator for a binding potential [33]:

$$iG(x,x') \;=\; iG^F(x,x') \;= < 0|T\{\psi(x)\bar{\psi}(x')\}|0> = \sum_{\alpha} u_{\alpha}(x)\bar{u}_{\alpha}(x')\theta(t-t') + \sum_{\beta} v_{\beta}(x)\bar{v}_{\beta}(x')\theta(t'-t) \tag{10}$$

Since the three-quark vacuum state $|0>$ does not contain any pion or gluon, the pion and gluon Green's functions are given by the usual free Feynman propagator for a boson field:

$$i\Delta_{ij}(x-x') \;= < 0|T\{\phi_i(x)\bar{\phi}_j(x')\}|0> = i\delta_{ij} \int \frac{d^4k}{(2\pi)^4} \frac{1}{k^2 - m_\pi^2 + i\epsilon}\, exp[-ik(x-x')]\,, \tag{11}$$

$$i\Delta_{ab}^{(\mu\nu)}(x-x') \;= < 0|T\{A_\mu^a(x)A_\nu^b(x')\}|0> = i\delta_{ab} g^{\mu\nu} \int \frac{d^4k}{(2\pi)^4} \frac{1}{k^2 + i\epsilon}\, exp[-ik(x-x')]\,, \tag{12}$$

(in the Coulomb gauge), where $g^{\mu\nu} = \delta_{\mu\nu} g^{\mu\mu}$, $g^{00} = -g^{11} = -g^{22} = -g^{33} = 1$.

On the basis of the effective Lagrangian and using the time-ordered perturbation theory within the frame of many-body quantum field theory [33] we can develop the calculation scheme for the excitation spectrum of the Nucleon and Delta. At zero-th order the quark core result (E_Q) is obtained by solving Eq.(4) for the single quark system numerically by using the harmonic oscillator basis. Since we work in the independent particle model, we assume that the bare three-quark state of the $SU(2)$-flavor baryons corresponds to the structure $(1S_{1/2})^2(nlj)$ with a single excited valence quark in the non-relativistic spectroscopic notation. Below, on the basis of the one-pion and one-gluon exchange mechanisms we will argue that such a configuration of the three valence quarks is identified with the baryon resonances decaying strongly into the $\pi + N$ channel. And contrary, the baryon states with more than one valence quarks in excited orbits do not have a strong coupling into this channel. In other words, all baryon resonances appearing in the πN scattering data can be identified with the above orbital configuration containing a single excited valence quark. This is why we fix the excited baryon configuration as $(1S_{1/2})^2(nlj)$. The corresponding quark core energy is evaluated as the sum of single quark energies with:

$$E_Q = 2E(1S_{1/2}) + E(nlj) \tag{13}$$

The second order perturbative corrections to the energy spectrum of the SU(2) baryons due to the pion ($\Delta E^{(\pi)}$) and gluon ($\Delta E^{(g)}$) fields are calculated on the basis of the Gell-Mann and Low theorem :

$$\Delta E = < \Phi_0| \sum_{i=1}^{\infty} \frac{(-i)^n}{n!} \int i\delta(t_1)\, d^4x_1 \ldots d^4x_n \, T[\mathcal{H}_I(x_1)\ldots\mathcal{H}_I(x_n)] \,|\Phi_0>_c \tag{14}$$

with $n = 2$, where the relevant quark-pion and quark-gluon interaction Hamiltonian densities are

$$\mathcal{H}_I^{(q\pi)}(x) = \frac{i}{f_\pi}\bar{\psi}(x)\gamma^5\vec{\tau}\vec{\phi}(x)S(r)\psi(x), \tag{15}$$

$$\mathcal{H}_I^{(qg)}(x) = g_s\bar{\psi}(x)A^a_\mu(x)\gamma^\mu\frac{\lambda^a}{2}\psi(x) \tag{16}$$

The stationary bare three-quark state $|\Phi_0>$ is constructed from the vacuum state using the usual creation operators:

$$|\Phi_0>_{\alpha\beta\gamma} = b^+_\alpha b^+_\beta b^+_\gamma|0>, \tag{17}$$

where α, β and γ represent the quantum numbers of the single quark states, which are coupled to the respective baryon configuration. The energy shift of Eq.(14) is evaluated up to second order in the quark-pion and quark-gluon interaction, and generates self-energy and exchange diagrams contributions. In the self-energy diagrams a single pion or gluon is emitted and absorbed by the same valence quark, which however can be excited to an intermediate quark or anti-quark state. In the second order exchange diagrams a single pion or gluon, emitted by a valence quark is absorbed by another valence quark of the SU(2) baryon.

B. Center of mass corrections

The result for E_Q in Eq.(13) contains an essential spurious contribution of the center of mass motion to the energy of the baryons. A covariant way of the separation of the CM motion is possible in non-relativistic models. In the nonrelativistic three nucleon system the energy is reduced by factor about 1/3 after the separation of the CM. At the same time different approaches are being used in relativistic mean field models.

For the ground state nucleon and delta baryons we use the development of the Ref. [34], where three different approximations have been used, which estimate corrections for the center of mass motion: the $R = 0$ [35], $P = 0$ [36] and LHO [37] methods. In all three methods the baryon wave function is rewritten in the Jacobi coordinates in the center of mass system as $\Phi_B(\vec{r}, \vec{\rho}, \vec{R})$, where $\vec{r}$, $\vec{\rho}$ and $\vec{R}$ are relative coordinates between the two valence quarks, between 3-valence quark and the center of mass of the 1+2 quarks, and the center of mass of the all three quarks, respectively.

In the R=0 method the baryon wave function in the CM system is multiplied by the plane wave of the CM motion:

$$\Phi_R(\vec{r_1}, \vec{r_2}, \vec{r_3}, \vec{P}) = N_R\ exp(i\vec{P}\cdot\vec{R})\Phi_B(\vec{r}, \vec{\rho}, \vec{R} = 0). \tag{18}$$

The second P=0 method is based on the Fourier transformation of the baryon wave function:

$$\Phi_P(\vec{r_1}, \vec{r_2}, \vec{r_3}, \vec{P}) = N_P\ exp(i\vec{P}\cdot\vec{R})\int exp(-i\vec{P}\cdot\vec{R'})\Phi_B(\vec{r}, \vec{\rho}, \vec{R'})d\vec{R'}. \tag{19}$$

The lowest harmonic oscillator (LHO) method is based on the projection of the baryon wave function on the lowest harmonic oscillator state:

$$\Phi_{LHO}(\vec{r_1}, \vec{r_2}, \vec{r_3}, \vec{P}) = N_{LHO}\ exp(i\vec{P}\cdot\vec{R})\int R_{0s}(\vec{R'})\Phi_B(\vec{r}, \vec{\rho}, \vec{R'})d\vec{R'}. \tag{20}$$

The factors N_R, N_P and N_{LHO} differ each from other and are found from the normalization conditions:

$$<\Phi(\vec{r_1}, \vec{r_2}, \vec{r_3}, \vec{P})|\Phi(\vec{r_1}, \vec{r_2}, \vec{r_3}, \vec{P'})> = (2\pi)^3\delta(\vec{P} - \vec{P'}). \tag{21}$$

In all of the three methods, the average kinetic energy and mass terms of the three-body system are estimated by using angular momentum algebra and numerical methods (see Ref.[34] for details).

For the excited nucleon and delta states we use a linear approximation based on the CM corrections of the ground state nucleon and delta. Similar estimation was used in the bag model [38]. This way is based on the statement that the CM correction does not affect any level ordering in the excited baryon spectrum and it must be at the same order for the both ground and excited baryon states [26]. The linear coefficients are found from the equation

$$k_{CM} = E_Q(CM)/E_Q, \tag{22}$$

where $E_Q(CM)$ is the CM-corrected quark core energy of the ground state nucleon, estimated by one of the above three methods, E_Q is the initial nucleon ground state quark core energy in Eq.(13). By using the same relation and using the value of the linear coefficient fixed from the ground state nucleon and delta, we can estimate the CM corrected quark core energy for the excited nucleon and delta states.

C. Self-energy diagrams contribution

The self-energy terms contain contribution both from intermediate quark ($E > 0$) and anti-quark ($E < 0$) states. These diagrams describe the processe when a pion or gluon is emitted and absorbed by the same valence quark which can be excited to the intermediate quark or anti-quark states.

The pion part of the self energy term (pion cloud contribution) (see Fig.1) is evaluated as

$$\Delta E^{(\pi)}_{s.e.} = -\frac{1}{2f_\pi^2}\sum_{a=1}^{3}\sum_{\alpha'\leq\alpha_F}\int\frac{d^3\vec{p}}{(2\pi)^3 p_0}\Big\{\sum_\alpha\frac{V^{a+}_{\alpha\alpha'}(\vec{p})V^{a}_{\alpha\alpha'}(\vec{p})}{E_\alpha - E_{\alpha'} + p_0} - \sum_\beta\frac{V^{a+}_{\beta\alpha'}(\vec{p})V^{a}_{\beta\alpha'}(\vec{p})}{E_\beta + E_{\alpha'} + p_0}\Big\}, \tag{23}$$

with $p_0^2 = \vec{p}^2 + m_\pi^2$. The $q - q - \pi$ transition form factors are defined as:

$$V^a_{\alpha\alpha'}(\vec{p}) = \int d^3x \bar{u}_\alpha(\vec{x})\Gamma^a(\vec{x})u_{\alpha'}(\vec{x})e^{-i\vec{p}\vec{x}} \tag{24}$$

$$V^a_{\beta\alpha'}(\vec{p}) = \int d^3x \bar{v}_\beta(\vec{x})\Gamma^a(\vec{x})u_{\alpha'}(\vec{x})e^{-i\vec{p}\vec{x}} \tag{25}$$

The vertex function of the $\pi - q - q$ and $\pi - q - \bar{q}$ transition is

$$\Gamma^a = S(r)\gamma^5\tau^a I_c\,, \tag{26}$$

where I_c is the color unity matrix. The expression of the $\pi - q - q$ transition form factor has been derived in Ref.[22]:

$$\begin{aligned} V^a_{\alpha\alpha'}(\vec{p}) \;=\; & \sum_{l_n}(-i)^{l_n+1}\int drr^2\Big[g_\alpha(r)f_{\alpha'}(r) + g_{\alpha'}(r)f_\alpha(r)\Big]S(r)j_{l_n}(pr) \\ & Y^{m'_j - m_j}_{l_n}(\hat{p})\mathcal{F}(l^\pm, l', l_n, j, j', m_j, m'_j) < m_t|\tau^a|m'_t >< m_c|I_c|m'_c > . \end{aligned} \tag{27}$$

The Hermitian conjunction of the transition form factor

$$\begin{aligned} V^{a+}_{\alpha\alpha'}(\vec{p}) \;=\; & \sum_{l_n}(i)^{l_n+1}\int drr^2\Big[g_\alpha(r)f_{\alpha'}(r) + g_{\alpha'}(r)f_\alpha(r)\Big]S(r)j_{l_n}(pr) \\ & Y^{(m'_j - m_j)*}_{l_n}(\hat{p})\mathcal{F}(l^\pm, l', l_n, j, j', m_j, m'_j) < m'_t|\tau^a|m_t >< m'_c|I_c|m_c > . \end{aligned} \tag{28}$$

After integration over the angular part in Eq. (17), the self-energy diagrams contribution to the baryon spectrum induced by pion fields is evaluated as:

$$\begin{aligned} \Delta E^{(\pi)}_{s.e.} = -\frac{1}{16\pi^3 f_\pi^2}\int\frac{dp\,p^2}{p_0}\sum_{\alpha'\leq\alpha_F}\sum_{l_n}\Big\{\sum_\alpha\frac{[\int drr^2 G_{\alpha\alpha'}(r)S(r)j_{l_n}(pr)]^2}{E_\alpha - E_{\alpha'} + p_0}Q_{s.e.}(l, l', l_n, j, j') - \\ \sum_\beta\frac{[\int drr^2 G_{\beta\alpha'}(r)S(r)j_{l_n}(pr)]^2}{E_\beta + E_{\alpha'} + p_0}Q_{s.e.}(l, l', l_n, j, j')\Big\}, \end{aligned} \tag{29}$$

where j_{l_n} is the Bessel function. The radial overlap of the single quark states with quantum numbers $\alpha = (N, l, j, m_j, m_t, m_c)$ and α' is defined as

$$G_{\alpha\alpha'}(r) = f_\alpha(r)g_{\alpha'}(r) + f_{\alpha'}(r)g_\alpha(r). \tag{30}$$

The angular momentum coefficients Q are evaluated for all SU(2) baryons as

$$Q_{s.e.}(l, l', l_n, j, j') = 12\pi[l^\pm][l_n][j]\Big[C^{l'0}_{l^\pm 0 l_n 0}W(j\tfrac{1}{2}l_n l'; l^\pm j')\Big]^2\sum_{m_j}\sum_{m'_j\leq\alpha_f}\Big[C^{j'm'_j}_{jm_j l_n(m'_j - m_j)}\Big]^2, \tag{31}$$

where C and W are the Clebsch-Gordan and Wigner coefficients, respectively.

The gluon part of the second order self-energy diagrams (gluon cloud) contribution is estimated in a similar way as

$$\Delta E^{(g)}_{s.e.} = \frac{g_s^2}{2}\sum_a g_{\mu\nu}\sum_{\alpha'\leq\alpha_F}\int\frac{d^3\vec{p}}{(2\pi)^3 p}\Big\{\sum_\alpha\frac{V^{a\mu+}_{\alpha\alpha'}(\vec{p})V^{a\nu}_{\alpha\alpha'}(\vec{p})}{E_\alpha - E_{\alpha'} + p} - \sum_\beta\frac{V^{a\mu+}_{\beta\alpha'}(\vec{p})V^{a\nu}_{\beta\alpha'}(\vec{p})}{E_\beta + E_{\alpha'} + p}\Big\}, \tag{32}$$



FIG. 1: Second order self energy diagrams induced by $\pi-$meson fields

where the transition form factor is evaluated with the corresponding vertex matrix

$$\Gamma^a_\mu = \gamma^\mu \frac{\lambda^a}{2} I_t \tag{33}$$

with the isospin unity matrix I_t.

$$V^{a\mu}_{\alpha\alpha'}(\vec{p}) = \delta_{\mu 0}\int d^3x \bar{u}_\alpha(\vec{x})\frac{\lambda_a}{2} I_t u_{\alpha'}(\vec{x}) exp(-i\vec{p}\vec{x}) + \delta_{\mu k}\int d^3x \bar{u}_\alpha(\vec{x})\frac{\lambda_a}{2} I_t \hat{\alpha}_k u_{\alpha'}(\vec{x}) exp(-i\vec{p}\vec{x}) \tag{34}$$

The last expression is convenient for the estimation of the exchange diagrams.

For the self-energy diagrams we use an alternative expression of the transition form-factors. Putting the quark wave functions with further integration over the radial part of the spatial coordinate one can write for the transition form-factor next equation:

$$V^{a\mu}_{\alpha\alpha'}(\vec{p}) = \sum_{l_n m_n}\sum_{LL'}\sum_{m_L m'_L m_s m'_s}\left(\frac{[L][l_n](4\pi)}{[L']}\right)^{\frac{1}{2}}(-i)^{l_n} Y_{l_n m_n}(\hat{p}) M^\mu_{m_s m'_s} C^{\mathrm{L'0}}_{\mathrm{L0l_n0}} C^{\mathrm{jm_j}}_{\mathrm{Lm_L\frac{1}{2}m_s}} C^{\mathrm{j'm'_j}}_{\mathrm{L'm'_L\frac{1}{2}m'_s}} C^{\mathrm{L'm'_L}}_{\mathrm{Lm_Ll_nm_n}} \cdot$$

$$\cdot \int r^2 R^{\alpha\alpha'}_{\mu_{\mathrm{LL'}}}(r) j_{\mathrm{l_n}}(pr) dr < m_t|I_t|m'_t >< m_c|\frac{\lambda_a}{2}|m'_c >, \tag{35}$$

where the spin transition matrices

$$M^0_{m_s m'_s} = \delta_{m_s m'_s},$$

and

$$M^k_{m_s m'_s} = \sum_{k'=\pm 1,0} h_{kk'}\left[\delta_{k'1}\delta_{m_s 1/2}\delta_{m'_s(-1/2)} + \delta_{k'(-1)}\delta_{m_s(-1/2)}\delta_{m'_s 1/2} + 2m_s\delta_{k'0}\delta_{m_s m'_s}\right]$$

with the only nonzero expansion coefficients $h_{1,+1} = h_{1,-1} = h_{3,0} = 1$, and $h_{2,+1} = -h_{2,-1} = -i$.

The radial functions are defined as

$$R^{\alpha\alpha'}_{\mu_{LL'}}(r) = \delta_{\mu,0}\delta_{Ll}\delta_{L'l'}(g_\alpha g_{\alpha'} + f_\alpha f_{\alpha'}) + i\delta_{\mu,k}(\delta_{Ll}\delta_{L'l'^{\pm}} g_\alpha f_{\alpha'} - \delta_{L'l'}\delta_{Ll^{\pm}} g_{\alpha'} f_\alpha)$$

The corresponding Feynman diagrams are given in Fig.2, where the contribution from intermediate quark and anti-quark levels have opposite signs.

After evaluation of the transition form-factors and integration over angular variables, the self-energy term induced by gluon fields can be written as a sum of color-electric (Coulomb) and color-magnetic parts (see Ref.[22]):

$$\Delta E^{(g)}_{s.e.} = \frac{g_s^2}{3\pi^2}\sum_{N'l'j'}\sum_{(\alpha,\beta)}\sum_{LL'L^*L'^*l_n}[l_n]\left(\frac{[L][L^*]}{[L'][L'^*]}\right)^{1/2} C^{L'0}_{L0l_n0} C^{L'^*0}_{L^*0l_n0}$$

$$\Big\{\delta_{lLL^*}\delta_{l'L'L'^*}\delta_{l_n l}\mathcal{A}^{jj'm_jm_j'}_{LL'L^*L'^*l_n}\left[\int\frac{[R_{\alpha\alpha'l_n}(p)+F_{\alpha\alpha'l_n}(p)]^2}{E_\alpha-E_{\alpha'}+p}pdp-\int\frac{[R_{\beta\alpha'l_n}(p)+F_{\beta\alpha'l_n}(p)]^2}{E_\beta+E_{\alpha'}+p}pdp\right]$$

$$-\left[\mathcal{B}^{jj'm_jm_j'}_{LL'L^*L'^*l_n}-\mathcal{D}^{jj'm_jm_j'}_{LL'L^*L'^*l_n}+2\mathcal{E}^{jj'm_jm_j'}_{LL'L^*L'^*l_n}\right]$$

$$\left[\int\frac{dp\,p}{E_\alpha-E_{\alpha'}+p}\mathcal{H}_{\alpha\alpha'l_nLL'L^*L'^*}-\int\frac{dp\,p}{E_\beta+E_{\alpha'}+p}\mathcal{H}_{\beta\alpha'l_nLL'L^*L'^*}\right]\Big\}, \tag{36}$$

where we define function

$$\mathcal{H}_{\alpha\alpha'l_nLL'L^*L'^*}=\mathcal{H}_{\alpha\alpha'l_nLL'L^*L'^*}(p)$$

$$=H^2_{\alpha\alpha'l_n}\delta_{lLL^*}\delta_{l'^\pm L'L'^*}+H^2_{\alpha'\alpha l_n}\delta_{l^\pm LL^*}\delta_{l'L'L'^*}-H_{\alpha\alpha'l_n}H_{\alpha'\alpha l_n}(\delta_{lL}\delta_{l^\pm L^*}\delta_{l'^\pm L'}\delta_{l'L'^*}+\delta_{lL^*}\delta_{l^\pm L}\delta_{l'L'}\delta_{l'^\pm L'^*}) \tag{37}$$

and radial integrals

$$H_{\alpha\alpha'l_n}=H_{\alpha\alpha'l_n}(p)=\int dr\left[r^2f_{\alpha'}(r)g_\alpha(r)j_{l_n}(pr)\right],$$

$$R_{\alpha\alpha'l_n}=R_{\alpha\alpha'l_n}(p)=\int dr\left[r^2g_{\alpha'}(r)g_\alpha(r)j_{l_n}(pr)\right],$$

$$F_{\alpha\alpha'l_n}=F_{\alpha\alpha'l_n}(p)=\int dr\left[r^2f_{\alpha'}(r)f_\alpha(r)j_{l_n}(pr)\right]. \tag{38}$$

The angular momentum coefficients $\mathcal{A}$, $\mathcal{B}$, $\mathcal{D}$ and $\mathcal{E}$ can be found from Appendix C of Ref. [22].



FIG. 2: Second order self energy diagrams induced by gluon fields

D. Exchange diagrams contribution

The pion exchange contribution to the baryon energy-shift (see Fig.3) is evaluated as:

$$\Delta E^{(\pi)}_{ex.}=-\frac{1}{2f_\pi^2}\sum_{a=1}^{3}\sum_{\alpha\le\alpha_F}\sum_{\alpha'\le\alpha_F}\int\frac{d^3\vec{p}}{(2\pi)^3p_0^2}\Big\{V^{a+}_{\alpha\alpha}(\vec{p})V^a_{\alpha'\alpha'}(\vec{p})-V^{a+}_{\alpha\alpha'}(\vec{p})V^a_{\alpha\alpha'}(\vec{p})\Big\}. \tag{39}$$

By using the Wick's theorem we can write a more convenient expression for the energy shift of the SU(2) baryons from the second order pion exchange diagrams:

$$\Delta E^{(\pi)}_{ex.} = -\frac{1}{16\pi^3 f_\pi^2}\int \frac{dp\,p^2}{p_0^2}\sum_{l_n}\Pi_{l_n}(p) \tag{40}$$

where

$$\Pi_{l_n}(p) =< \Phi_B|\sum_{i\neq j}\vec{\tau}(i)\vec{\tau}(j)T_{ln}(i)T_{l_n}(j)K_{l_n}(i)K^+_{l_n}(j)|\Phi_B > \tag{41}$$

and the operators $\vec{\tau}, T_{l_n}$ and K_{l_n} are summed over single quark levels $i \neq j$ of the SU(2) baryon. In the quark model, the baryon wave function $|\Phi_B>$ is presented as a bound state of three valence quarks in the orbital configuration $(1S)^2(nlj)$, and it can be written down commonly as

$$|\Phi_B >= |\alpha\beta\gamma > (J_0T_0) = |\alpha\beta;\gamma >^{TM_T(T_0)}_{JM(J_0)}$$
$$= \hat{S}\Big[|\psi_\alpha(r_1)\psi_\beta(r_2)\psi_\gamma(r_3)\mathcal{Y}^{JM}_{J_0}(\hat{x_1}\hat{x_2};\hat{x_3}) > |\chi^{TM_T}_{T_0}(12;3) >\Big]|\chi^{(}_c 123) >,$$

where J_0 and T_0 are intermediate spin and isospin couplings of the two S-wave valence quarks, respectively. They satisfy the symmetry requirement $S_0 = T_0$. The states ψ are the single particle states, labeled by a set of quantum numbers α, β and γ, excluding the color degree of freedom.



FIG. 3: Second order $\pi-$meson exchange diagrams

The operator T_{l_n} in equation (41) is the radial integration operator:

$$< \alpha|T_{l_n}|\beta >= \int dr\Big[r^2 S(r) j_{l_n}(pr) G_{\alpha\beta}(r)\Big]. \tag{42}$$

with

$$G_{\alpha\alpha'}(r) = f_\alpha(r)g_{\alpha'}(r) + f_{\alpha'}(r)g_\alpha(r). \tag{43}$$

where $\alpha = (N, l, j, m_j, m_t, m_c)$ and α' are two sets of the single quark quantum numbers. The matrix elements of the operator K_{l_n} are given by

$$< \alpha|K_{l_n}|\beta >= -\Big(4\pi[l^{\pm}(\alpha)][l_n][j(\alpha)]\Big)^{1/2} C^{l(\beta)0}_{l^{\pm}(\alpha)0l_n0}$$
$$W(j(\alpha)\frac{1}{2}l_n, l(\beta); l^{\pm(\alpha)}, j(\beta))C^{j(\beta)m(\beta)}_{j(\alpha)m_j(\alpha)l_n(m(\beta)-m(\alpha))}, \tag{44}$$

and the Hermitian conjunction

$$< \alpha|K^{+}_{l_n}|\beta > = < \beta|K_{l_n}|\alpha >,$$

where $j(\alpha), l(\alpha), l^{\pm}(\alpha), m(\alpha)$ are the quantum numbers of the single quark state $< \alpha|$.

The contribution of the second-order gluon-exchange terms to the baryon spectrum (see Fig.4) is given by

$$\Delta E^{(g)}_{ex.} = -\frac{g^2}{2}\sum_{a\mu\nu}\sum_{\alpha\le\alpha_F}\sum_{\alpha'\le\alpha_F}\int\frac{d^3\vec{p}}{(2\pi)^3p^2}\Big\{V^{a\mu+}_{\alpha\alpha}(\vec{p})V^{a\nu}_{\alpha'\alpha'}(\vec{p}) - V^{a\mu+}_{\alpha\alpha'}(\vec{p})V^{a\nu}_{\alpha\alpha'}(\vec{p})\Big\}g^{\mu\nu}. \tag{45}$$

By using the Wick's theorem we can write more convenient expression for this equation

$$\Delta E^{(g)}_{ex.} = -\frac{g^2}{\pi}\int\limits_0^{\infty} dp \sum_{l_n m_n}\mathcal{Q}_{l_n m_n}(p) \tag{46}$$

with the corresponding color-electric (Coulomb) and color-magnetic parts:

$$\begin{aligned}\mathcal{Q}_{l_n m_n}(p) = &< \Phi_B|\sum_{i\neq j}\frac{\vec{\lambda}(i)}{2}\frac{\vec{\lambda}(j)}{2}T^{(g)}_{ln}(i)T^{(g)}_{ln}(j)\hat{F}_{l_n m_n}(i)\hat{F}^{+}_{l_n m_n}(j)|\Phi_B > \\ &- < \Phi_B|\sum_{i\neq j}\frac{\vec{\lambda}(i)}{2}\frac{\vec{\lambda}(j)}{2}T^{(g)}_{ln}(i)T^{(g)}_{ln}(j)\hat{F}_{l_n m_n}(i)\hat{F}^{+}_{l_n m_n}(j)\hat{\vec{\alpha}}(i)\hat{\vec{\alpha}}(j)|\Phi_B > .\end{aligned} \tag{47}$$

The operator $T^{(g)}_{ln}$ is the radial integration operator with the factor $j_{l_n}(pr)$. The operators $\hat{F}_{l_n m_n}(i)$ and $\hat{F}^{+}_{l_n m_n}(j)$ are the angular integration operator with the factors $Y_{l_n m_n}(\hat{x}_i)$ and $Y^{*}_{l_n m_n}(\hat{x}_j)$ respectively. All these operators are summed over single quark levels $i \neq j$ of the SU(2) baryon.



FIG. 4: Second order gluon-exchange diagrams

E. Selection rules for the quantum numbers of the excited N^* and Δ^* states

Now we begin to analyze the excited N^* and Δ^* spectrum based on the relativistic description of one-pion and one-gluon (color magnetic part) exchange mechanisms. These exchange operators, as was found in Ref.[22], couple the upper and lower components of the two interacting valence quarks, respectively. Based on this fact we can derive the selection rules for the quantum numbers of the baryon states with the fixed orbital configuration.

Let us to fix the orbital configuration as $(1S_{1/2})^2(nlj)$, with the intermediate spin coupling $\vec{S}_0 = \vec{S}_1 + \vec{S}_2 = \vec{1/2} + \vec{1/2}$ of the two $1S$-valence quarks, where the last valence quark (nlj) can be in the ground or an excited state. The upper and lower Dirac components of the last excited valence quark have orbital momenta l and $l' = l \pm 1$, respectively. Our choice of the above orbital configuration is close to the limitation in the diquark-quark models [39], where some of the degrees of freedom are "frozen". The corresponding baryon states are different from members of the SU(6)⊗O(3) multiplets in the Constituent Quark Models.

The first two selection rules come from the coupling of the three valence quarks into the SU(2) baryon state with total momentum J and isospin T:

$$\begin{aligned} \vec{S}_0 + \vec{j} &= \vec{J} \\ \vec{S}_0 + \vec{1/2} &= \vec{T}, \end{aligned} \tag{48}$$

where the symmetry property of the two S-quark coupling was used. The third rule comes from the pion exchange mechanism between the excited valence quark and the $1S$ quark. This mechanism couples the upper (lower) component of the $1S$ valence quark with the lower (upper) component of the excited (nlj) valence quark. Since the upper component of the S-quark has zero orbital momentum, then for the orbital momentum of the exchanged pion we derive the equation

$$L_\pi = l' = l \pm 1 \tag{49}$$

The final selection rule is based on the assumption that the coupling of the last valence quark with quantum numbers (nlj) to the $1S$ quark plus pion is the main component of the strong coupling of the excited baryon state to the $N(939) + \pi$:

$$\vec{L}_\pi + \vec{1/2} = \vec{J} \tag{50}$$

With this assumption, Eq.(49) can be used for the identification of the baryon resonance in the πN-scattering process. Namely, when $l' = 0$ we have S-wave nucleon and delta resonances, when $l' = 1$ we have P-wave resonances, etc.

An important consequence of the obtained selection rules is that all the N^* and Δ^* resonances appearing in the πN scattering process and coupling strongly to the πN channel are identified with the orbital configurations $(1S_{1/2})^2(nlj)$ with two valence quarks in the ground state and a single valence quark in an excited state. A baryon resonance corresponding to the orbital configuration with two valence quarks in excited states $(1S_{1/2})(nlj)_1(nlj)_2$ couples strongly to the $\pi\pi N$-channel, but not to the πN channel.

Using the obtained selection rules it is very natural to analyze schematically the excited nucleon and delta spectrum. For the fixed orbital configurations $(1S_{1/2})^2(nlj)$ with the intermediate spin coupling of the two $S-$ wave quarks $S_0 = 0$ (the so-called instanton channel), Eq.(48) allows only a single N^* state with $J = j$ and no any Δ^* resonances.

Except the case, when the last valence quark is in the $P_{1/2}$ orbit, the intermediate coupling $S_0 = 1$, due-to the selection rule Eq.(50) yields two resonances in the both nucleon and delta sectors with the total momentum $J = L_\pi \pm 1/2$. In this way one of the N^* resonances defined by the selection rules in Eq.(48) with $J = j + 1$ or $J = j - 1$ is ruled out. When the last valence quark is in the $P_{1/2}$ orbit, i.e. has the lower S-component, the selection rules yield $L_\pi = 0$ and $J = 1/2$, and consequently, only single S-wave resonances in the both nucleon and Δ sectors are allowed.

Thus, for the fixed $(1S_{1/2})^2(nlj)$ orbital configuration with $(nlj) \neq (nP_{1/2})$ there must be a band of three N^* and two Δ^* resonances. The lightest N^* state corresponds to the intermediate spin coupling $S_0 = 0$ due-to strong attraction in this "instanton channel". The other two N^*, as well as the two Δ^* resonances correspond to the spin coupling $S_0 = 1$ and must be close each to other.

In the case when the last quark is in the $P_{1/2}$ orbit, there is a band of two N^* states (not close each to other) and a single Δ^* resonance appearing in the S-wave of the πN scattering.

III. NUMERICAL RESULTS

A. Condition of the calculations

In order to account for the finite size effect of the pion, we introduce a one-pion vertex regularization function in the momentum space, parameterized in the dipole form as

$$F_\pi(p^2) = \frac{\Lambda_\pi^2 - m_\pi^2}{\Lambda_\pi^2 + p^2}.$$

We fix $\Lambda_\pi = 1$ GeV in our calculations. Contrary to the bag-model calculations, the above regularization is used not for the solution of the convergence problem of the quark self-energy. This was explicitly shown in Ref.[25] and [23] for the lowest valence quark states. As is known from Ref.[18], the convergence of the quark self-energy is a serious problem in the bag models.

As was noted above, the strength $c = 0.16$ GeV^2 and Coulomb $\alpha = \pi/12 \approx 0.26$ parameters of the Cornell potential are fixed from the flux-tube study [32] and lattice calculations [27, 28]. However, it is useful to note that the above value of the strength parameter was already probed long times ago in Ref.[26]. The only free parameter of the model, m of the confining potential was chosen as m=60 MeV to reproduce the correct axial charge of the proton $g_A = 1.26$ (and the empirical pion-nucleon coupling constant $G^2_{\pi NN}/4\pi = 14$ via the Goldberger-Treiman relation). It yields a reasonable value for the quark core RMS radius of the proton 0.52 fm (see [26]). The strong coupling constant $g_s^2 = 4\pi\alpha_s$ with the value $\alpha_s = 0.65$.

In Ref. [26] by examining the different model parameters the sensitivity of the Nucleon energy on the description of the static properties of the proton has been examined. It was found that a larger value of the strength parameter c of the confining potential yields a smaller value for the proton RMS radius.

Stating that the Coulomb like term of the Cornell potential $V(r) = -\alpha/r$ is actually due-to the color electric component of one-gluon exchange mechanism, we need to avoid a double counting of these components in the calculations of gluon loop corrections to the baryon mass spectrum. This is why we have restricted our study to the color-magnetic component of the one-gluon exchange forces together with one-pion loop corrections.

In Ref.[23] we have demonstrated explicitly a convergence of the self-energy for the valence quarks in the lowest $1S$, $2S$, $1P_{1/2}$, $1P_{3/2}$ orbits induced by the pion and color-magnetic gluon fields. The total momentum of the intermediate quark and anti-quark states increases from $j = 1/2$ up to $j = 25/2$, while their radial quantum numbers grows up to $n = 20$ in order to reach convergent results.

We have obtained convergence of the self-energy also for the excited valence quark states in the orbits $3S$, $2P_{1/2}$, $2P_{3/2}$, $1D_{5/2}$, which are included into the structure $(1S)^2(nlj)$ of the excited baryons in present study.

By summing the self energies of the three valence quarks in the excited $(1S)^2(nlj)$ nucleon and Delta states, we can estimate the contribution of the self-energy terms to the excitation spectrum of the SU(2) flavor baryons.

B. Ground state Nucleon spectrum

In Table 1 we give the mass values for the g.s. N(939) with and without CM correction in three different methods: the R=0, [35], P=0 [36] and LHO [37]. All these methods were firstly examined in Ref. [34]. As we can see from the Table, they agree within 50 MeV for the ground state Nucleon.

The pion loop diagrams yield positive contribution to the baryon mass-spectrum due-to self-energy term. For the ground state Nucleon it is 200 MeV.

For the gluon field contributions we probe two different ways. In the first case we include the contribution of all the intermediate quark and antiquark states up to convergence with j =25/2. The corresponding results are given in the 3-row of Table 1, they increases the Nucleon mass by 109 MeV. In the second case a restriction of the intermediate states to the ground $1S$ quark state is used when estimating the self-energy (I=0). The second approximation is based on the short-range character of one-gluon exchange forces. The corresponding energy shift for the ground state Nucleon is now negative (-127 MeV). However, after including the center of mass corrections, the Nucleon mass is still overestimated by about 100 MeV. In principle, we can fit the strong coupling constant α_s to reproduce the N(939) mass value, but first we have to check the excitation spectrum of the SU(2) flavor baryons. Thus, from the results in Table 1 we can conclude, that the second way, when the short-range character of one-gluon exchange forces is taken into account, is most favorable.

We note that the agreement within 50 MeV of the three R=0 [35], P=0 [36] and LHO [37] methods for the CM correction is reasonable. Moreover, these three methods always give corrections with systematic differences. Namely, the LHO method always yields correction larger than the $P = 0$ method, but smaller than the $R = 0$ method (see Ref. [22]). The values of the linear coefficient for the CM-correction in Eq.(22) for these three methods are 0.548 (R=0), 0.574 (P=0) and 0.563 (LHO). Thus, we can fix one of these methods (LHO) and go to the excited sector.

C. Spectrum of the SU(2) flavor baryons

In Table 2 we compare our numerical estimations of the excited N^* and Δ^* spectrum within the developed schematic periodic table with the last experimental data from [6] and [7]. The calculations were done up-to and including F-wave baryon resonances in the frame of the developed chiral quark model. In the Table we give the center of mass (CM) corrected quark core results (zero order estimation) (second column) together with the second order pion field contributions corresponding to the self energy (3-th column) and exchange diagrams (4-th column). The 5-th column contains results for the quark core plus pion loop corrections. Next 6-th and 7-th columns correspond to the contributions of the self-energy and exchange terms of the color-magnetic one-loop diagrams. The final theoretical estimations are given in the 8-column with the strong coupling constant $\alpha_s = 0.65$. As was argued above, due to the short range character of the gluon exchange forces between valence quarks, we restrict our calculations of the color-magnetic self-energy terms to the case, where the intermediate quark is the same initial and final quark.

Based on obtained selection rules first we will show the assignment of the excited baryon states presented in the data from Ref.[6] with corresponding orbital configurations. Let us to fix the orbital configuration $(1S_{1/2})^2(nS_{1/2})$. In the data there are four N^* with $J^\pi = 1/2^+$ (P_{11} resonances) and two N^* with $J^\pi = 3/2^+$ (P_{13} resonances). With the above rules, we can find easily that $N^*(1440)$, $N^*(1710)$, and $N^*(1720)$ resonances belong to the orbital configuration $(1S_{1/2})^2(2S_{1/2})$ with the radially excited $2S$ valence quark state, while the other three $N^*(1880)$, $N^*(1900)$ and $N^*(2100)$ resonances correspond to the orbital configuration $(1S_{1/2})^2(3S_{1/2})$. In the Δ sector there are two resonances with $J^\pi = 3/2^+$ at 1600 MeV and 1920 MeV, and two states with $J^\pi = 1/2^+$ at 1750 MeV and 1910 MeV which belong to the orbital configuration with the radially excited valence quark in consistence with our results.

The orbital configuration $(1S_{1/2})^2(1D_{3/2})$ is not presented in the data, since it would give two N^* resonances with $J^\pi = 3/2^+$ and a single N^* resonance with $J^\pi = 1/2^+$.

For the orbital configurations $(1S_{1/2})^2(nP_{1/2})$ there are four nucleon and three delta resonances with $J^\pi = 1/2^-$ and they are not close each to others. Each of the nucleon bands $n = 1$ and $n = 2$ contains two resonances, while Δ^* resonances correspond to the three bands including $n = 3$.

The orbital configuration $(1S_{1/2})^2(nP_{3/2})$ with $n = 1$ yields three N^* resonances $3/2^-(1520)$, $5/2^-(1675)$ and $3/2^-(1700)$, the first of which is less than other two states in accordance with our prediction. The band with $n = 2$ yields next group of the D-wave Nucleon resonances $3/2^-(1860)$, $3/2^-(2080)$ and $5/2^-(2200)$.

In the Delta sector there are four D-wave resonances, however only two of them $\Delta(5/2^-)(1930)$ and $\Delta(3/2^-)(1940)$ are close each to other. Since other D-wave resonances $\Delta(3/2^-)(1700)$ and $\Delta(5/2^-)(2350)$ are far, then we can predict possible new $\Delta^*(5/2^-)$ (around 1700 MeV) and $\Delta^*(3/2^-)$ (around 2350 MeV) resonances.

The F-wave N^* resonances $N^*(5/2^+)(1680)$, $N^*(5/2^+)(1870)$ and $N^*(7/2^+)(1990)$ belong to the orbital configuration $(1S_{1/2})^2(nD_{5/2})$ with $n = 1$ together with delta states $\Delta^*(5/2^+)(1905)$ and $\Delta^*(7/2^+)(1950)$, while the $\Delta^*(5/2^+)(2000)$ and $\Delta^*(7/2^+)(2390)$ belong to the $n = 2$ band.

We can continue our analysis at higher energies and predict in summary seven new N^* resonances with $J^\pi = 7/2^-$ (2000 MeV), $9/2^+$ (2100 - 2300 MeV), $11/2^+$ (2100 - 2300 MeV), $11/2^-$ (2500-2700 MeV), $13/2^-$ (2500-2700 MeV), $13/2^+$ (2600 -2800 MeV), $15/2^+$ (2600 -2800 MeV) and four Δ^* resonances with $J^\pi = 5/2^-$ (around 1700 MeV), $3/2^-$ (2350 MeV), $11/2^-$ (2750 MeV), $13/2^+$ (2950 MeV). These resonances are expected to be observed in current experimental facilities.

It is clear now that the remaining "missing N^* and Δ^* resonances" predicted by the Constituent Quark Models must appear in the $\pi\pi N$ strong coupling sector, if they exist. As we have argued above, they will be assigned with the orbital configuration $(1S_{1/2})(nlj)_1(nlj)_2$ with two excited valence quarks and a single ground state valence quark.

Now we can analyze the numerical values within our model in comparison with the experimental data from Ref. [6]. In Fig.5 and Fig.6 we give the theoretical estimations and experimental data in a convenient diagrammatic way. Table 2 contains information about orbital configurations for each baryon resonance, as well as separate contributions from self-energy and exchange diagrams due-to pion- and color-magnetic gluon fields. As can be seen from the Table and figures, the mass spectrum of the Nucleon and Δ is described reasonably well in the relativistic chiral quark model with a single free parameter of the confining potential.

For the test of the results we can check the consistence of our results with the results of the Cloudy Bag Model [16]. The pion exchange diagrams contribute about 144 MeV to the energy difference between $N(939)$ and $\Delta(1232)$, while the gluon exchange forces yield 64 MeV for the strong coupling constant value $\alpha_s = 0.65$. The value $\alpha_s = 1.51$ increases the gluon field contribution up to 149 MeV, which is consistent with the CBM results. However, as one can see from the Table, this way strongly moves down almost all the baryon states including $N(939)$ and $\Delta(1232)$.

The next important observation is that one needs an additional exchange mechanism for the lowering both the ground state $N(939)$, Roper resonance $N(1440)$ and $N^*(1900)(3/2^+)$ almost by the same amount. On the other hand, four of the radially excited nucleon resonances: $N^*(1710)(1/2^+)$, $N^*(1880)(1/2^+)$, $N^*(1720)(3/2^+)$ and $N^*(2100)(1/2^+)$ are inside the corresponding error boxes.

The close situation is in the Δ sector. A state $\Delta(1750)1/2^+$ is described well, while other three radially excited resonances together with $\Delta(1212)$ (slightly) are overestimated.

Contrary, the orbitally excited N^* resonances are mostly underestimated. The situation in the Δ^* sector is different. The orbitally excited Δ^* states corresponding to the lowest radial quanta $n = 1$ are underestimated, while negative parity Δ states corresponding to the radial quantum number n=2 are consistent with the experimental data.

It is relevant to compare the obtained estimations for the excited N* an d $\Delta*$ spectrum with the results of the relativized Constituent Quark Model [3]. A comparison of the results presented in the Fig.5 and Fig.6 with the results presented in Fig.9 and Fig.10 of above mentioned work indicates that the two methods describe the excited baryon spectrum at the same level. However, the present model does not have any fitting parameters, and , additionally, unlike CQM, it does not predict many missing nonobserved states.

The analysis shows that one needs an additional exchange mechanism between valence quarks to reproduce the whole SU(2) baryon spectrum. The new exchange forces must depend on the spin and flavor of valence quarks as well as on the quantum numbers of the baryon state. Of course, large part of the interaction comes from two-pion exchange mechanism.



FIG. 5: Spectrum of the Nucleon states. Theoretical estimations (solid lines) in comparison with experimental data (boxes) from Ref.[6]

IV. CONCLUSIONS

In summary, we have derived selection rules for the excited baryon state, assuming that it's orbital configuration is of the form $(1S)^2(nlj)$ with two valence quarks in the ground state and a single excited quark. These selection rules were derived on the basis of the one-pion exchange mechanism between valence quarks in the frame of the relativistic chiral quark model. An important consequence of the obtained selection rules is that all the N^* and Δ^* resonances appearing in the πN scattering process and strongly coupling to the πN channel are identified with the orbital configurations $(1S_{1/2})^2(nlj)$. Baryon resonances corresponding to the orbital configuration with two valence quarks in excited states couple strongly to the $\pi\pi N$-channel, but not to the πN channel.



FIG. 6: Spectrum of the Delta states (notations are the same as in Fig.5)

Based on obtained selection rules, we have constructed a schematic periodic table and calculate the energy spectrum of the excited N^* and Δ^* baryons within the field-theoretical framework including one-pion and one-gluon loop corrections. The obtained numerical estimations for the energy positions of baryon resonances (up to and including F-wave) yield an overall good description of the experimental data. However, Nucleon ground state and some of the radially excited baryon resonances (including Roper) are overestimated. Contrary, all the orbitally excited N^* and a number of the Δ^* resonances are underestimated. At the same time, the remaining Nucleon and Delta states including $\Delta(1232)$ are well reproduced.

The important observation is that one needs an additional exchange mechanism for the lowering both the ground state $N(939)$ and Roper resonance $N(1440)$ almost by the same amount. This fact indicate that the "lowering mechanism" for the both N(939) and Roper resonance $N^*(1440)(1/2^+)$ should be the same, since these states have identical quantum numbers except the radial quantum number. Of course, the two-pion exchange forces are expected to contribute essentially to the excited baryon spectrum.

A comparison of the obtained results with the results of the relativized Constituent Quark Model indicates that they describe the excited baryon spectrum at the same level. This level of description in our model was achieved without any fitting parameters. Moreover, unlike CQM, our model does not yield many nonobserved resonances at the lower excitation spectrum. The only $\Delta(5/2^-)$ resonance is expected to be observed at energy scale around 1600-1800 MeV.

At higher energies, where the experimental data are poor, we can extend our model schematically and predict seven new N^* and four Δ^* states with larger spin values. Of course, the number of "missing resonances" in our model is strongly suppressed due-to restriction of the configuration space to the orbits $(1S_{1/2})^2(nlj)$. However, as we have shown above, at lower energies this construction works reasonably well.

Acknowledgments

Author thanks Th. Gutsche for his valuable advices and discussions, S. Krewald and A. Rakhimov for useful discussions, and K. Shimizu for the help in the calculations of the center-of mass corrections.

[1] L.Y. Glozman, W. Plessas, K. Varga and R.F. Wagenbrunn, Phys. Rev. **D58**, 094030 (1998).
[2] S. Capstick and N. Isgur, Phys. Rev. **D34**, 2809 (1986)
[3] S. Capstick and W. Roberts, PPNP, **V45**, 241 (2000).
[4] U. Löring, K. Kretzschmar, B.C. Metsch and H.R. Petry, Euro.Phys.J. **A10**, 309 (2001);
U. Löring, B.C. Metsch and H.R. Petry, Euro.Phys.J. **A10**, 395 (2001);
U. Löring, B.C. Metsch and H.R. Petry, Euro.Phys.J. **A10**, 447 (2001);
B. Metsch, U. Löring, D. Merten and H. Petry, EPJ **A18**, 189 (2003).
[5] K. Nakamura et. al., J. Phys. **G37**, 075021 (2010).
[6] E. Klempt and J.M. Richard, Rev. Mod. Phys. **82**, 1095 (2010).
[7] A.V. Anisovich, E. Klempt, V.A. Nikonov, A.V. Sarantsev and U. Thoma, arXiv:1009.4803 (2010).
[8] K. Sasaki and S. Sasaki, Phys. Rev. **D72**, 034502 (2005);
K. Sasaki, S. Sasaki and T. Hatsuda, Phys. Lett. **B623**, 208 (2005)
[9] N. Mathur, Y. Chen, S.J. Dong, T. Draper, I. Horvath, F.X. Lee, K.F. Liu, J.B. Zhang, Phys. Lett. **B605**, 137 (2005).
[10] H.L.L. Roberts, I. LeiChang, C. Cloet and C.D. Roberts, Few Body Syst. **51**, 1 (2011).
[11] I.T. Obukhovsky, A. Faessler, D.K. Fedorov, T. Gutsche and V.E. Lyubovitskij, Phys. Rev. **D84**, 014004 (2011).
[12] D. Muller, M. Buballa and J. Wambach, Phys.Rev. **D81**, 094022 (2010).
[13] J. Luecker, C.S. Fischer and R. Williams, Phys.Rev. **D81**, 094005 (2010).
[14] O. Krehl, C. Hanhart, S. Krewald and J. Speth, Phys.Rev. **C62**, 025207 (2000);
M. Doering, C. Hanhart, F. Huang, S. Krewald and U.-G. Meissner, Nucl.Phys. **A829**, 170 (2009);
M. Doering, C. Hanhart, F. Huang, S. Krewald, U.-G. Meissner and D. Roenchen, Nucl.Phys. **A851**, 58 (2011).
[15] H. Kamano and T.-S.H. Lee, arXiv:1108:0324 (2011).
[16] S. Theberge, A.W. Thomas and G.A. Miller, Phys. Rev. **D22**, 2838 (1980);
A.W. Thomas, S. Theberge and G.A. Miller, Phys. Rev. **D24**, 216 (1981).
[17] A.W. Thomas, Prog.Part.Nucl.Phys. **61**, 219 (2008);
F. Myhrer and A.W. Thomas, Phys.Lett. **B663**, 302 (2008).
[18] K. Saito, Prog. Theor. Phys. **V71**, 775 (1984).
[19] S. Nagai, T. Miyatsu and K. Saito, Phys. Lett. **B666**, 239 (2008).
[20] M. Wakamatsu, Eur.Phys. J. **A44**, 297 (2010).
[21] E.M. Tursunov, J. Phys. G: Nucl. Part. Phys. **31**, 617 (2005).
[22] E.M. Tursunov, J. Phys. G: Nucl. Part. Phys. **36**, 095006 (2009).
[23] E.M. Tursunov, J. Phys. G: Nucl. Part. Phys. **37**, 105013 (2010).
[24] E. Oset, R. Tegen and W. Weise, Nucl. Phys. A **426**, 456 (1984).
[25] Th. Gutsche and D. Robson, Phys. Lett. **B229**, 333 (1989).
[26] Th. Gutsche "A Chiral Potential Model for the Light Quark Baryons", Ph.D thesis, Florida State Univ., 1987, 249 p.
[27] T. Kawanai and S. Sasaki, PPNP, **67**, 130 (2012); arXiv:1110.0888 (2011).
[28] J. Greensite, and S. Olejnik, Phys.Rev. **D67**, 094503 (2003).
[29] M. Gell-Mann and M. Levy, Nuovo Cim. **16**, 1729 (1960).
[30] W. Weise, Inter. Rev. of Nucl. Phys., Vol.1, ed. W. Weise (World Scientific, Singapore), p.58, 1984.
[31] S. Coleman, "Soft Pions", in: Hadrons and Their Interactions, Acad. Press, 1968
[32] M. Lüscher, Nucl. Phys. **B180**, 317 (1981).
[33] A.I. Fetter and J.D. Waleska, "Quantum theory of many particle systems" (McGraw-Hill, New York), 1971.
[34] Y.B. Dong, K. Shimizu, A. Faessler and A.J. Buchmann, Phys.Rev. **C60**, 035203 (1999).
[35] D.H. Lu, A.W. Thomas and A.G. Williams, Phys.Rev. **C57**, 2628 (1998).
[36] R. Tegen, R. Brockmann and W. Weise, Z.Phys. **A307**, 339 (1982).
[37] L. Wilets, "Non-Topological Solitons (World Scientific, Singapoure), 1989.
[38] T.A. DeGrand, Ann. of Phys. **100**, 425 (1976).
[39] M. Anselmino, E. Predazzi, S. Ekelin, S. Fredriksson and D.B. Lichtenberg, Rev. Mod. Phys. **65**, 1199 (1993).

TABLE I: The mass value of the g.s. nucleon in MeV with and without center of mass (CM) correction

	No CM	R=0, [35]	P=0, [36]	LHO, [37]
E_Q	1715	940	985	966
$E_Q + \Delta E(\pi)$	1915	1140	1185	1166
$E_Q + \Delta E(\pi + g)$	2024	1249	1294	1275
$E_Q + \Delta E(\pi + g)$, I=0	1788	1013	1058	1039

TABLE II: Estimations for the energy values of the N^* and Δ^* resonances in MeV

SU(2) baryon state	$E_Q(CMcor.)$	$\Delta E_\pi^{s.e.}$	$\Delta E_\pi^{ex.}$	$E_Q + \Delta E_\pi$	$\Delta E_g^{s.e.}$	$\Delta E_g^{ex.}$	E(theor)	E(exp.)[6]
$N(939)(1/2^+)(P_{11})\ (1S)^3$	966	380	-180	1166	-95	-32	1039	938 ÷ 939
$N(1440)(1/2^+)(P_{11})\ (1S)^2(2S)$	1200	603	-113	1690	-70	-24	1596	1430 ÷ 1470
$N(1710)(1/2^+)(P_{11})\ (1S)^2(2S)$	1200	603	-66	1737	-70	-10	1657	1650 ÷ 1750
$N(1720)(3/2^+)(P_{13})\ (1S)^2(2S)$	1200	603	1	1804	-70	10	1744	1700 ÷ 1760
$N(1880)(1/2^+)(P_{11})\ (1S)^2(3S)$	1361	788	-110	2039	-66	-28	1945	1840 ÷ 1940
$N(2100)(1/2^+)(P_{11})\ (1S)^2(3S)$	1361	788	-39	2110	-66	-1	2043	2000 ÷ 2200
$N(1900)(3/2^+)(P_{13})\ (1S)^2(3S)$	1361	788	-3	2146	-66	11	2089	1900 ÷ 2000
$N(1535)(1/2^-)(S_{11})\ (1S)^2 1P_{1/2}$	1129	501	-119	1511	-79	-13	1419	1528÷1548
$N(1650)(1/2^-)(S_{11})\ (1S)^2 1P_{1/2}$	1129	501	46	1676	-79	-49	1548	1640÷1680
$N(1905)(1/2^-)(S_{11})\ (1S)^2 2P_{1/2}$	1301	713	-111	1903	-69	-27	1807	1850÷1950
$N(2090)(1/2^-)(S_{11})\ (1S)^2 2P_{1/2}$	1301	713	24	2038	-69	-13	1956	2100÷2260
$N(1520)(3/2^-)(D_{13})\ (1S)^2 1P_{3/2}$	1107	515	-126	1496	-91	-27	1378	1518 ÷ 1526
$N(1700)(3/2^-)(D_{13})\ (1S)^2 1P_{3/2}$	1107	515	-79	1543	-91	-9	1443	1675 ÷ 1775
$N(1675)(5/2^-)(D_{15})\ (1S)^2 1P_{3/2}$	1107	515	11	1633	-91	29	1571	1670 ÷ 1680
$N(1860)(3/2^-)(D_{13})\ (1S)^2 2P_{3/2}$	1293	713	-111	1895	-73	-29	1793	1810 ÷ 1890
$N(2080)(3/2^-)(D_{13})\ (1S)^2 2P_{3/2}$	1293	713	-31	1975	-73	-1	1901	2045 ÷ 2155
$N(2200)(5/2^-)(D_{15})\ (1S)^2 2P_{3/2}$	1293	713	4	2010	-73	20	1957	2075 ÷ 2245
$N(1680)(5/2^+)(F_{15})\ (1S)^2 1D_{5/2}$	1212	638	-114	1736	-89	-30	1617	1680 ÷ 1690
$N(1870)(5/2^+)(F_{15})\ (1S)^2 1D_{5/2}$	1212	638	-37	1813	-89	2	1726	1840 ÷ 1960
$N(1990)(7/2^+)(F_{17})\ (1S)^2 1D_{5/2}$	1212	638	12	1862	-89	27	1800	1860 ÷ 2100
$\Delta(1232)(3/2^+)(P_{33})\ (1S)^3$	966	380	-36	1310	-95	32	1247	1230 ÷ 1234
$\Delta(1600)(3/2^+)(P_{33})\ (1S)^2(2S)$	1200	603	-23	1780	-70	34	1744	1535 ÷ 1695
$\Delta(1750)(1/2^+)(P_{31})\ (1S)^2(2S)$	1200	603	1	1804	-70	8	1742	1710 ÷ 1780
$\Delta(1910)(1/2^+)(P_{31})\ (1S)^2(3S)$	1361	788	-3	2146	-66	23	2103	1845 ÷ 2025
$\Delta(1920)(3/2^+)(P_{33})\ (1S)^2(3S)$	1361	788	-18	2131	-66	8	2073	1880 ÷ 2020
$\Delta(1620)(1/2^-)(S_{31})\ (1S)^2 1P_{1/2}$	1129	501	-24	1606	-79	45	1572	1603 ÷ 1649
$\Delta(1900)(1/2^-)(S_{31})\ (1S)^2 2P_{1/2}$	1301	713	-24	1990	-69	12	1933	1860 ÷ 1960
$\Delta(1700)(3/2^-)(D_{33})\ (1S)^2 1P_{3/2}$	1107	515	-18	1604	-91	8	1521	1670 ÷ 1770
$\Delta(5/2^-)(D_{35})\ (1S)^2 1P_{3/2}$	1107	515	-35	1587	-91	35	1531	...
$\Delta(1940)(3/2^-)(D_{33})\ (1S)^2 2P_{3/2}$	1293	713	-9	1997	-73	9	1933	1935 ÷ 2055
$\Delta(1930)(5/2^-)(D_{35})\ (1S)^2 2P_{3/2}$	1293	713	-22	1984	-73	28	1939	1900÷ 1960
$\Delta(1905)(5/2^+)(F_{35})\ (1S)^2 1D_{5/2}$	1212	638	-12	1838	-89	7	1756	1860 ÷ 1940
$\Delta(1950)(7/2^+)(F_{37})\ (1S)^2 1D_{5/2}$	1212	638	-27	1823	-89	29	1763	1915 ÷ 1960